

Eastern Health Collaborative Quality Measures

Measure	Corresponding National Measures	Description	Numerator Statement (measurement period is the last 18 months unless otherwise noted)	Denominator Statement (measurement period is the last 18 months unless otherwise noted)	Goal
Comprehensive Diabetes Care: HbA1c Poor Control (>9.0%)	NQF 0059 PQRS 1	The percentage of patients 18-75 years of age with diabetes (type 1 and 2) whose most recent HbA1c level during the measurement year was greater than 9.0% (poor control) or was missing a result, or if an HbA1c test was not done during the measurement year.	Patients whose most recent HbA1c level (performed during the measurement period) is >9.0%	Patients 18-75 years of age with diabetes with a visit during the measurement period	Less than 40% of patients have A1c \geq 9
Comprehensive Diabetes Care: Eye Exam	NQF 0055 PQRS 117	The percentage of patients 18-75 years of age with diabetes (type 1 and 2) who had an eye exam (retinal) performed.	Patients with an eye screening for diabetic retinal disease. This includes diabetics who had one of the following: A retinal or dilated eye exam by an eye care professional or a negative retinal exam (no evidence of retinopathy) by an eye care professional	Patients 18-75 years of age with diabetes with a visit during the measurement period	At least 60% of diabetic patients have a yearly eye exam
Comprehensive Diabetes Care: Influenza Vaccines		The percentage of patients 18-75 years of age with diabetes (type 1 and 2) and are adequately immunized with influenza vaccine.	Diabetic patients who received an influenza immunization OR who reported previous receipt of an influenza immunization	All diabetic patients aged 18-75 seen for a visit between October 1 and March 31	80% of diabetic patients have received an influenza vaccine
Comprehensive Diabetes Care: Pneumonia Vaccines		The percentage of patients 18-75 years of age with diabetes (type 1 and 2) and are adequately immunized with pneumonia vaccine.	Diabetic patients who have ever received a pneumococcal vaccination	All diabetic patients 18-75 years of age	80% of diabetic patients have received a pneumonia vaccine
Preventive Care Screening: Tobacco Use: Screening and Cessation	NQF 0028	Percent of patients aged 18 years and older who were screened for tobacco use at least once during the two-year measurement period AND who received cessation counseling interventions if identified as a tobacco user.	Patients who were screened for tobacco use at least once within 24 months AND who received tobacco cessation counseling intervention if identified as a tobacco user	All patients aged 18 years and older	50% of Tobacco users be offered a cessation program

Eastern Health Collaborative Quality Measures

<p>Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up</p>	<p>NQF 0421 PQRS 128</p>	<p>Percentage of patients aged 18 years and older with a calculated BMI in the past six months or during the current visit documented in the medical record AND if the most recent BMI is outside of normal parameters, a follow-up plan is documented.</p> <p style="text-align: center;">Normal Parameters: Age 65 years and older BMI \geq 23 and $<$30 Age 18 – 64 years BMI \geq 18.5 and $<$25</p>	<p>Patients with a documented BMI during the encounter or during the previous six months, AND when the BMI is outside of normal parameters, a follow-up plan is documented during the encounter or during the previous six months of the encounter with the BMI outside of normal parameters.</p>	<p>All patients 18-75 years of age</p>	<p>50% have BMI recorded in EMR and have documented counseling for weight loss</p>
<p>Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents</p>	<p>NQF 0024 PQRS 239</p>	<p>Percentage of children 3-17 years of age who had an outpatient visit with a primary care physician (PCP) or an OB/GYN and who had evidence of body mass index (BMI) percentile documentation, counseling for nutrition and counseling for physical activity during the measurement year.</p>	<p>Numerator 1: Patients who had a height, weight and body mass index (BMI) percentile recorded during the measurement period and who had counseling for nutrition during the measurement period and who had counseling for physical activity during the measurement period</p>	<p>Patients 3-17 years of age with at least one outpatient visit with a primary care physician (PCP) or an obstetrician/ gynecologist (OB/GYN) during the measurement period</p>	<p>50% have BMI recorded in EMR and have documented counseling for weight loss</p>