



# Clinical Quality Measures Workgroup

## Proposed Change to Project Measures 4 of 16

Compiled for the IHC 4/8/2016 V1.0

Measure	Original Measure Language	Proposed Measure Language	CQM Comments
1*	Percentage of patients identified as tobacco users who received cessation intervention during the two-year measurement period.	Percentage of patients aged 18 years and older who were screened for tobacco use one or more times within 24 months AND who received cessation counseling intervention if identified as a tobacco user.	Aligns with CMS 138, PQRS 226, NQF 0028.
2*	Percentage of children, two through 17 years of age, whose weight is classified based on Body Mass Index (BMI), who receive counseling for nutrition and physical activity.	Percentage of patients 3-17 years of age who had an outpatient visit with a Primary Care Physician (PCP)** or Obstetrician/Gynecologist (OB/GYN) and who had evidence of the following during the measurement period. Three rates are reported. <ul style="list-style-type: none"> <li>- Percentage of patients with height, weight, and body mass index (BMI) percentile documentation</li> <li>- Percentage of patients with counseling for nutrition</li> <li>- Percentage of patients with counseling for physical activity</li> </ul>	Aligns with CMS 155, PQRS 239, NQF 0024.  ** This measure is includes all providers such as Physicians, Physician Assistants, and Nurse Practitioners in Family Medicine, Primary Care Medicine, Internal Medicine, General Practice Medicine, Pediatric Medicine, or Obstetrician/Gynecologist (OB/GYN) Medicine. The data is collected based on procedures conducted rather than the type of provider.
3*	The percentage of patients 18-75 with a diagnosis of diabetes, who have optimally managed modifiable risk factors (A1c<8.0%, LDL<100 mg/dL, blood pressure<140/90 mm Hg, tobacco non-use, and daily aspirin usage for patients with diagnosis of IVD with the intent of preventing or reducing future complications associated with poorly managed diabetes.	Percentage of patients 18-75 years of age with diabetes who had hemoglobin A1c > 9.0% during the measurement period	Aligns with CMS 122, PQRS 001, NQF 0059.  There are other national measures for diabetes but most present collection challenges. Others have not been updated to reflect the latest clinical best practices.
4	The percentage of members 18 to 74 years of age who had an outpatient visit and who's BMI was documented during the measurement year or the year prior to the measurement year.	Percentage of patients aged 18 years and older with a BMI documented during the current encounter or during the previous six months AND with a BMI outside of normal parameters, a follow-up plan is documented during the encounter or during the previous six months of the current encounter.	Aligns with CMS 69, PQRS 128, NQF 421.

\* These are CMMI Year 1 required measures; states may select a national measure or similar state specific metric.

\*\* This measure is inclusive of all providers including Physicians, Physician Assistants, and Nurse Practitioners in Family Medicine, Primary Care Medicine, Internal Medicine, General Practice Medicine, Pediatric Medicine, or Obstetrician/Gynecologist (OB/GYN) Medicine. The data is collected based on procedures conducted rather than the type of provider.



Statewide Healthcare Innovation Plan (SHIP) is supported by Funding Opportunity Number CMS-1G1-14-001 from the U.S. Department of Health and Human Services, Center for Medicare & Medicaid Services.